WHAT IS CLAIMED IS:

Sub C'> 1. An isolated nucleic acid specific to Histoplasma capsulatum comprising:

a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1;

a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1;

a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1;

a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1;

a fragment of a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1;

a fragment of a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1;

a fragment of a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1; or

a fragment of a nucleic acid having a nucleotide sequence which is substantially the same as a nucleic acid which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1;

where in the nucleic acid does not contain the nucleotide sequences 5 CGAAGTCGAGGCTTTCAGCATG3, 5 TATTAGCTCTAGAATTACCACGGGTATCCAAGTAGTAAGG3, 5 CCCCGAAGGGCATTGGTTTTTTATCTAATAAATACACCCC3, or nucleotide sequences complementary thereto,

and wherein the nucleic acid is not a nucleic acid

10

1 C C C 20

25

30

consisting essentially of between 10 and 100 nucleotides which is able to form a hybrid at 60°C with a nucleotide polymer having a nucleotide sequence of 5 CGAAGTCGAGGCTTCAGCATG3, 5 CATGCTGAAAGCCTCGACTTCG3, 5 CAUGCUGAAAGCCUCGACUUCG3 or 5 CGAAGUCGAGGCUUUCAGCAUG3.

5

10

CO

√0 ∭ 15

LA C. L. T. C. L.

20

25

- 2. The nucleic acid of Claim 1, wherein the nucleic acid has the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 3. The nucleic acid of Claim 1, wherein the nucleic acid has a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 4. The nucleic acid of Claim 1 wherein the nucleic acid has a nucleotide sequence which is substantially the same as a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1.
- 5. The nucleic acid of Claim 1, wherein the nucleic acid has a nucleotide sequence which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 6. The nucleic acid of Claim 1, wherein the nucleic acid is a fragment of a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1.
- 7. The nucleic acid of Claim 1, wherein the nucleic acid is a fragment of a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.

25

30

5

- 8. The nucleic acid of Claim 1, wherein the nucleic acid is a fragment of a nucleic acid having a nucleotide sequence which is substantially the same as a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1.
- 9. The nucleic acid of Claim 1, wherein the nucleic acid is a fragment of a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEO ID NO:1.
- 10. An isolated or recombinantly-produced antigen specific to *Histoplasma capsulatum* comprising:
- a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1;
- a fragment of a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1;
- a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1; or
- a fragment of a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 11. The antigen of Claim 10, wherein the antigen is a polypeptide encoded by a nucleic acid having a

nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.

12. The antigen of Claim 11, wherein the antigen has an amino acid sequence as set forth in the Sequence Listing as SEQ ID NO:2.

10

15

20

30

TOTAL TOTAL

- 13. The antigen of Claim 10, wherein the antigen is a fragment of a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 14. The antigen of Claim 10, wherein the antigen is a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 15. The antigen of Claim 10, wherein the antigen is a fragment of a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 16. A vector comprising a nucleic acid specific to 25 Histoplasma capsulatum, wherein the nucleic acid:

has a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1;

has a nucleotide sequence which is substantially the same as a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1;

is a fragment of a nucleic acid having a nucleotide

5

sequence as set forth in the Sequence Listing as SEQ ID NO:1; or

is a fragment of a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1; and wherein the vector is suitable for expressing the nucleic acid.

- 17. The vector of Claim 16, wherein the nucleic acid has a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1.
 - 18. The vector of Claim 16, wherein the nucleic acid has a nucleotide sequence which is substantially the same as a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1.
 - 19. The vector of Claim 16, wherein the nucleic acid is a fragment of a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1.
- 20. The vector of Claim 16, wherein the nucleic acid is a fragment of a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1.

10

15

20

25

30

35

- 21. A method for detecting a previous or current Histoplasma capsulatum infection in a subject, comprising:
- (a) contacting a fluid or tissue sample from the subject which contains antibodies with an isolated or recombinantly-produced antigen which is specific to Histoplasma capsulatum; and
- (b) detecting the presence of binding between the antibodies and the antigen, the presence of binding indicating the presence of a previous or current Histoplasma capsulatum infection in a subject,

wherein the antigen is:

a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1;

a fragment of a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1;

a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1; or

a fragment of a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.

22. The method of Claim 21, wherein the antigen is a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1.

- a fragment of a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 25. The method of Claim 21, wherein the antigen is a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.

- 26. The method of Claim 21, wherein the antigen is a fragment of a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
 - 27. An isolated antibody produced against an antigen of Claim 10.
 - 28. A kit for detecting a previous or current Histoplasma capsulatum infection in a sample comprising:
 - (a) an isolated nucleic acid of Claim 1, an isolated or recombinantly-produced antigen of Claim 10 or an isolated antibody of Claim 27; and
- (b) instructions describing the use of the nucleic acid, antigen or antibody in the detection of a previous
 30 or current Histoplasma capsulatum infection.

10

15

20

- 29. The kit of Claim 28, wherein the kit contains an antigen, and the antigen is a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 30. The kit of Claim 28, wherein the kit contains an antigen, and the antigen is a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 31. A host for expressing a polypeptide specific to Histoplasma capsulatum comprising a vector containing a nucleic acid, wherein the vector is suitable for expressing the nucleic acid, and wherein the nucleic acid:

has a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1;

has a nucleotide sequence which is substantially the same as a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1;

is a fragment of a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1; or

- is a fragment of a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1.
- 32. The host of claim 31, wherein the nucleic acid has a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1.

10

15

20

- 33. The host of Claim 31, wherein the nucleic acid is substantially the same as a nucleic acid having a nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1.
- 5 34. A method for detecting a past exposure to the fungus *Histoplasma capsulatum* comprising:
 - (a) injecting intradermally in the skin of a patient an effective amount of an isolated or recombinantly-produced antigen which is specific to Histoplasma capsulatum; and
 - (b) observing the skin at the injection site at a predetermined time after injection for a presence of swelling of the skin, the presence of swelling of the skin indicating a past exposure by the patient to the fungus Histoplasma capsulatum, wherein the polypeptide is:
 - a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1;

an antigenic fragment of a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1;

- a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1; or
- an antigenic fragment of a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.

35. The method of Claim 34, wherein the antigen is a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence as set forth in the Sequence Listing as SEQ ID NO:1.

5

- 36. The method of Claim 34, wherein the antigen is an antigenic fragment of a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 37. The method of Claim 34, wherein the antigen is a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 38. The method of Claim 34, wherein the antigen is a fragment of a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.
- 39. A vaccine for the prevention of histoplasmosis comprising:
- 25 (a) an effective amount of a nucleic acid of claim
 1 or an isolated or recombinantly-produced antigen of
 Claim 10; and
 - (b) a pharmaceutically-acceptable carrier.

10

15

20

25

30

- 40. A method for detecting a current *H. capsulatum* infection in a subject suspected of having an *H. capsulatum* infection comprising:
- (a) contacting a fluid or tissue sample from the subject which contains antigens with antibodies generated against an antigen which contains an epitope which is unique to *H. capsulatum*; and
- (b) detecting the presence of binding between the antigens and the antibodies, the presence of binding indicating the presence of a current *H. capsulatum* infection in the subject,

wherein the antigen is:

a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1;

a fragment of a polypeptide encoded by a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1;

a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1; or

a fragment of a polypeptide encoded by a nucleic acid which is substantially the same as a nucleic acid having a nucleotide sequence which is complementary to the nucleotide sequence set forth in the Sequence Listing as SEQ ID NO:1.

